

STRUCTURE FUNCTIONS ARE NOT PARTON PROBABILITIES

S. Peigné^a, F. Sannino^b

^aLAPTH, BP 110, F-74941 Annecy-le-Vieux Cedex, France

^bNORDITA, Blegdamsvej 17, DK-2100 Copenhagen, Denmark

We explain why contrary to common belief, the deep inelastic scattering structure functions are not related to parton probabilities in the target.

(Talk presented during the ‘International Light-Cone Workshop’, Trento, ECT*, September 3-11, 2001.)

Our statement originates from noting the incompatibility between the Glauber-Gribov picture of nuclear shadowing and the deep inelastic scattering (DIS) cross section σ_{DIS} being determined by parton probabilities. In section 1 we present physical arguments for this incompatibility. In section 2 we prove our main statement by an explicit calculation in a scalar QED model.

Our talk is based on the work done in [1], to which we refer for technical details.

1. PARTON DISTRIBUTIONS VERSUS GLAUBER-GRIBOV NUCLEAR SHADOWING

According to the QCD factorization theorem [2], σ_{DIS} can be expressed, at leading-twist and in the Bjorken limit*, as a convolution between hard partonic subprocess cross sections and parton distributions. The quark distribution in the nucleon N of momentum p reads

$$f_{q/N}(x_B, Q^2) = \int \frac{dy^-}{8\pi} e^{-ix_B p^+ y^-} \times \langle N | \bar{q}(y^-) \gamma^+ P \exp \left[ig \int_0^{y^-} dw^- A^+(w^-) \right] q(0) | N \rangle \quad (1)$$

Note that the (frame-independent) expression for $f_{q/N}$ is written in (1) with a particular choice of

*Choosing $q = (\nu, \vec{0}_\perp, q^z)$ and $p = (M, \vec{0}_\perp, 0)$ for the momenta of the virtual photon and of the target, the Bjorken limit corresponds to $Q^2 = -q^2 \rightarrow \infty$ and $\nu \rightarrow \infty$ with $x_B = Q^2/(2M\nu) = Q^2/M(q^- + q^+)$ being fixed. We use the light-cone variables $q^\pm = q^0 \pm q^z$.

Lorentz frame, namely $q^z < 0$. In the Bjorken limit we then have $q^- \simeq 2\nu \gg q^+$ and $q^+ = -Q^2/q^- \simeq -Mx_B < 0$. In such a $q^+ < 0$ frame, the virtual photon fluctuation $\gamma^* \rightarrow q\bar{q}$ is forbidden in light-cone time-ordered (LCTO) perturbation theory. Hence only LCTO diagrams where γ^* is *absorbed* can contribute. In this framework initial (ISI) and final (FSI) state interactions are then defined as occurring before or after the γ^* absorption time.

Gauge invariance of the matrix element in (1) is ensured by the presence of the path-ordered exponential. The latter is responsible, in Feynman gauge and more generally in covariant gauges, for FSI of the struck quark p_1 (see Fig. 1) which affect σ_{DIS} . The path-ordered exponential reduces to the identity in the light-cone $A^+ = 0$ gauge. Thus if (1) is correct also in this particular gauge, as usually believed, $f_{q/N}$ is then given by the square of the target nucleon wave function (evaluated in $A^+ = 0$ gauge) [3]. This would allow a probabilistic interpretation of the structure functions. In other words, only ISI would contribute to σ_{DIS} in $A^+ = 0$ gauge.

We will demonstrate in section 2, in the framework of a simple model, that FSI do modify σ_{DIS} in *all* gauges, including the $A^+ = 0$ gauge. Hence $f_{q/N}$ cannot be simply interpreted as a parton probability and its expression (1), although valid in covariant gauges, does not actually hold in $A^+ = 0$ gauge. The fact that this gauge is plagued with singularities has already been pointed out in [2]. Our results suggest that a more general expression for $f_{q/N}$ taking into ac-

count the $A_\perp$ components of the target gauge field is needed. These components might formally be included by writing the path-ordered exponential in the covariant form $P \exp [ig \int_0^y dw_\mu A^\mu(w)]$.

The Glauber-Gribov picture of nuclear shadowing [4,5] illustrates why structure functions and parton probabilities cannot be simply related. Roughly speaking the shadowing-type contribution is encoded in the structure functions and arises as a quantum interference effect between (in general *complex*) rescattering amplitudes while parton probabilities are related only to the *real* wave function of a stable target.

More specifically in a Lorentz frame where $q^+ > 0$, the $\gamma^* \rightarrow q\bar{q}$ fluctuation is possible in a LCTO formulation and occurs long before the target for $x_B \ll 1$. An interference can arise between an amplitude where the $q\bar{q}$ pair scatters inelastically on a target nucleon N_2 and another where it undergoes first an elastic scattering (via Pomeron exchange) on a nucleon N_1 (see Fig. 1). At small x_B the interference is destructive because the exchanged Pomeron amplitude becomes purely imaginary while the intermediate state after the elastic scattering is quasi on-shell. But note that the presence of on-shell intermediate states (OSIS) is independent of the Lorentz frame *and* of the gauge. Hence the Glauber-Gribov mechanism viewed in the frame of interest $q^+ < 0$ (where the quark distribution (1) is expressed) and in $A^+ = 0$ gauge must also involve OSIS. These OSIS cannot arise from ISI (they would constitute decay channels of the target) but from FSI. Since shadowing affects σ_{DIS} , this suggests that FSI modify σ_{DIS} (*i.e.* do not cancel when summing over cuts in the forward DIS amplitude) in all gauges including $A^+ = 0$. This contradicts the probabilistic interpretation of parton distributions, according to which in the $A^+ = 0$ gauge only ISI, which build the wave function, can influence the DIS cross section.

2. A PERTURBATIVE SCALAR QED MODEL FOR RESCATTERING

We now present a toy model which captures the main features of the shadowing phenomenon. Since the color and spin degrees of freedom are




Figure 1. Glauber-Gribov shadowing involves interference between rescattering amplitudes.

not essential for our conclusions, we consider scalar QED and compute the specific contribution to the forward DIS amplitude depicted in Fig. 2. As target T , we take a scalar ‘heavy quark’ of mass M . The incoming virtual photon couples with charge e to a scalar ‘light quark’ of mass m . The heavy and light quarks interact via ‘gluon’ (photon) exchanges with coupling g . As we will see, the particular contribution of Fig. 2 involving four ‘gluon’ exchanges exhibits the features of shadowing mentioned in section 1. The square of the Pomeron exchange amplitude is obtained when cutting the diagram between the exchanges k_2 and k_3 while cutting between k_3 and k_4 gives the interference of Fig. 1 (two of the three gluons of the left amplitude then model a Pomeron). The contribution of the diagram shown in Fig. 2 to σ_{DIS} is negative (see (6)) and thus reduces the Born cross section (corresponding to two exchanged gluons instead of four).

Let us note that since we consider an elementary target, our model is not meant to really describe nuclear shadowing, but is however adequate to study the effect of FSI on the cross section. Our perturbative treatment allows to study precisely the structure of what would be called the ‘soft’ part of the dynamics in QCD factorization theorems [2]. We take the hard part at zeroth order in g and focus on the aligned jet kinematics (where the ‘hard’ quark p_1 takes all the incoming photon energy whereas the antiquark p_2 remains ‘soft’) in the Bjorken limit. We then consider the small x_B limit both for simplicity and to recover features of shadowing. Our final statements how-




Figure 2. Forward $\gamma^* T \rightarrow \gamma^* T$ amplitude. All attachments of the exchanged gluons to the upper scalar loop are included, as well as topologically distinct permutations of the lower vertices on the target line.

ever do not depend on the small x_B limit. We consider the kinematics:

$$2\nu \sim p_1^- \gg p_2^- \gg k_{i\perp}, p_{i\perp}, k_i^-, m \gg k_i^+ \quad (2)$$

In order to evaluate the sum of the three relevant cuts of the forward $\gamma^*(q)T(p) \rightarrow \gamma^*(q)T(p)$ amplitude, we need to calculate three basic ingredients, the amplitudes $\gamma^*(q)T(p) \rightarrow q(p_1)\bar{q}(p_2)T(p')$ corresponding to one-, two- and three-gluon exchange which we call A , B and C respectively. Typical diagrams contributing to B in the small x_B limit and in Feynman gauge are shown in Fig. 3 (see [1] for more details).

These amplitudes can be expressed in the transverse coordinate space ($\vec{r}_\perp$ denotes the transverse separation between the quark p_1 and antiquark p_2 , and $\vec{R}_\perp$ the distance between the target quark and the light $q\bar{q}$ pair) [1]:

$$\begin{aligned} A &= 2eg^2MQp_2^- V(m_{||}r_\perp)W(\vec{r}_\perp, \vec{R}_\perp) \\ B &= \frac{-ig^2}{2!}WA \\ C &= \frac{(-ig^2)^2}{3!}W^2A \end{aligned} \quad (3)$$

where

$$\begin{aligned} m_{||}^2 &= p_2^- Mx_B + m^2 \\ V(mr_\perp) &= \frac{1}{2\pi}K_0(mr_\perp) \\ W(\vec{r}_\perp, \vec{R}_\perp) &= \frac{1}{2\pi} \log \left(\frac{|\vec{R}_\perp + \vec{r}_\perp|}{R_\perp} \right) \end{aligned} \quad (4)$$

The function V denotes the incoming longitudinal photon wave function. In the case of scalar quarks, the longitudinal photon polarization indeed dominates in the Bjorken limit. The function W stands for the amplitude associated to dipole scattering. Thus $A \propto W$, $B \propto W^2$ and $C \propto W^3$. Our results (3) are gauge independent[†]. Note that B is imaginary in the small x_B limit and $B \propto p_2^-$, hence B incorporates the main features of Pomeron exchange.




Figure 3. Diagrams which give leading order contributions to the one-loop amplitude B in Feynman gauge.

The contribution of Fig. 2 to σ_{DIS} is

$$\Delta \left(Q^4 \frac{d\sigma}{dQ^2 dx_B} \right) = \frac{\alpha}{16\pi^2} \frac{1-y}{y^2} \frac{1}{2M\nu} \int \frac{dp_2^-}{p_2^-} \times \int d^2\vec{r}_\perp d^2\vec{R}_\perp [|B|^2 + 2AC] \quad (5)$$

This is easily seen to be non-zero and negative:

$$|B|^2 + 2AC = -\frac{1}{3} \frac{g^4}{4} A^2 W^2 < 0 \quad (6)$$

so that (5) gives a negative correction to the Born DIS cross section given by A^2 , as expected for shadowing. The interference term $\sim 2AC$ overcompensates the $|B|^2$ term.

The rescattering correction to σ_{DIS} is leading-twist since $|B|^2 + 2AC \propto Q^2$, which makes the r.h.s. of (5) independent of Q in the Bjorken limit.

In Feynman gauge it can be shown that the partial contribution to (5) from rescatterings of the

[†]We obtained (3) in both Feynman and $A^+ = 0$ gauges. In $A^+ = 0$ gauge, we showed the result to be independent of the prescription used to regularize the spurious $k_i^+ = 0$ poles [1].

‘soft’ antiquark p_2 vanishes[‡]. Thus only rescatterings of the hard quark p_1 contribute to (6) in Feynman gauge. This is nothing else than the confirmation of the presence of FSI in covariant gauges.

Now comes our central argument. At small x_B the two-gluon exchange amplitude B $\gamma^*(q)T(p) \rightarrow q(p_1)\bar{q}(p_2)T(p')$ arises from the intermediate state occurring between the first and second/last gluon exchange being on-shell. (This intermediate state becomes exactly on-shell in the $x_B \rightarrow 0$ limit.) The typical LC distance y^- between the exchanges behaves as $\sim 1/x_B \rightarrow \infty$. This can be seen by calculating B in LCTO perturbation theory, where every intermediate state is associated with a denominator factor

$$D_{int} = \sum_{inc} k^- - \sum_{int} k^- + i\varepsilon \quad (7)$$

which measures the LC energy difference between the incoming and intermediate states. An intermediate state is on-shell when only $\sum_{inc} k^- - \sum_{int} k^- = 0$ contributes to the amplitude. In the matrix element (1), all rescatterings occur at equal time y^+ , whereas the LC distance $y^- \sim 1/x_B$. Both our amplitudes B and C arise from such OSIS in the $x_B \rightarrow 0$ limit. (An infinite distance is allowed between any of the gluon exchanges.) Obviously, the expression $|B|^2 + 2AC$, built from B and C , also arises purely from OSIS. The presence of OSIS being gauge independent, we conclude that B and C , and thus $|B|^2 + 2AC$, originate from OSIS also in $A^+ = 0$ gauge. As stated previously, these OSIS cannot arise from ISI for a stable target. They can be created only via FSI. We conclude:

In any gauge FSI are present and do modify the DIS cross section. Thus structure functions cannot be simply related to parton probabilities.

Acknowledgements. It is a pleasure for us to thank S. Brodsky, P. Hoyer and N. Marchal for a very stimulating collaboration.

REFERENCES

1. S. Brodsky, P. Hoyer, N. Marchal, S. Peigné and F. Sannino, hep-ph/0104291, submitted to Phys.Rev.D.
2. J. C. Collins and D. E. Soper, Nucl. Phys. **B194**, 445 (1982);
J. C. Collins, D. E. Soper and G. Sterman, Nucl. Phys. **B261**, 104 (1985), Nucl. Phys. **B308**, 833 (1988), Phys. Lett. **B438**, 184 (1998) and review in *Perturbative Quantum Chromodynamics*, (A.H. Mueller, ed., World Scientific Publ., 1989, pp. 1-91);
G. T. Bodwin, Phys. Rev. **D31**, 2616 (1985), Erratum Phys. Rev. **D34**, 3932 (1986).
3. S. J. Brodsky, M. Diehl and D. S. Hwang, Nucl. Phys. **B596**, 99 (2001) [hep-ph/0009254];
M. Diehl, T. Feldmann, R. Jakob and P. Kroll, Nucl. Phys. **B596**, 33 (2001) [hep-ph/0009255].
4. V. N. Gribov, Sov. Phys. JETP **29**, 483 (1969) and **30**, 709 (1970).
5. G. Piller and W. Weise, Phys. Rep. **330**, 1 (2000) [hep-ph/9908230];
G. Piller, M. Vanttinen, L. Mankiewicz and W. Weise, hep-ph/0010037.

[‡]This is also true in a general covariant gauge, but not in $A^+ = 0$ gauge. Indeed the subset of diagrams involving only rescatterings of p_2 is gauge dependent.